



Effect of spawn density and different bed substrate on the sporophore production of *Lentinus edodes* (Berkley)

C.KANNAN

Department of Plant Pathology, Annamalai University.

Abstract

L. edodes is largely cultivated in China, Japan and other Asian countries because of its taste and nutritional values. It is important nutritionally because of its higher protein, dietary fibers and important mineral contents. The present study was investigated the various spawn density and different bed substrate on the sporophore production of *Lentinus edode*. The revealed that that the spawn density of 20 per cent level completely colonized the beds within 31 days and the bed substrates, silver oak saw dust recorded shortest spawn run days.

Introduction

Shiitake is globally a well known cultivated species, but yet to find a place in Indian markets. Lack of cultivation technology on locally available substrates and suitable high temperature strains are the reason for its non-availability in India (Sharma *et al.*, 2006). Despite sporadic research efforts to standardize its cultivation technology in India, it could not reach the commercial level, as available technology is not viable. Hence, in view of the potentials of Shiitake to find a place in Indian markets the present research has been carried out with various spawn density and different bed substrate on the sporophore production of *Lentinus edodes*

Materials and methods

Effect of spawn density on the sporophore yield of *Lentinus edodes*

The ill filled paddy grain spawn was prepared and inoculated into the bed substrate @ 5, 10, 15, 20, 25 and 30 per cent levels on w/w basis following layer spawning method (Sivaprakasam, 1980). The beds were prepared with different levels of spawn densities and incubated at $25\pm 2^{\circ}\text{C}$. The observations on spawn run, pin head formation, number of sporophores, yield and biological efficiency were assessed and recorded following standard procedures.

Efficacy of bed substrates on the growth and yield of *Lentinus edodes*



Nine different locally available substrates *viz.*, paddy straw, sugarcane trash, sugarcane baggase, coir pith, groundnut shell, various saw dusts *viz.*, silver oak, teak, eucalyptus and coconut were evaluated for their efficacy in enhancing the yield of *L.edodes*.

Substrate preparation

Paddy straw, Sugarcane trash and Sugarcane bagasse

Well dried chopped substrates *viz.*, Paddy straw, Sugarcane trash and Sugarcane bagasse @ 1 kg / bed, previously soaked in water for 12 hours, followed by steam sterilization (15 psi for 30 min.) were used for the preparation of cylindrical (polybag 60×30cm) beds similar to that of oyster mushroom bed preparation (Thiribhuvanamala *et al.*, 2005 a). At the time of bed preparation the moisture level of the substrates was maintained at 50 per cent.

Saw dust

The method enumerated by Royse (2005) and was followed for bed preparation. The various sawdust substrates obtained from *viz.*, silver oak, teak, eucalyptus and coconut were steam sterilized (15 psi for 30 min.) and used for the preparation of beds as described earlier. At the time of bed preparation, the moisture level of the saw dust was maintained at 50 per cent by assessing in such a way that upon squeezing no water trickles down the substrate but the palm is wetted.

Coir pith

Partially decomposed coir pith was steam sterilized (15 psi for 30 min.) and used for the preparation of beds as described earlier.

Groundnut shells

Well dried groundnut shells crushed into small bits were steam sterilized (15 psi for 30 min.) and used for the preparation of beds as described earlier. After the preparation the beds were pressed gently and tied with polythene strip. The IFPG spawn of *L.edodes* was inoculated to the beds at 20 per cent level following layer spawning method. A temp. of 25±2°C and relatively humidity 90 % were maintained in the cropping room. After spawn running, when the browning process nearing completion, the polypropylene bag was removed and the beds were dipped in cold water (5°C) for ten min. so as to induce basidiocarp



formation. Fully matured basidiocarps were harvested periodically and the parameters like number of days taken for spawn run, days taken for first harvest, total yield and biological efficiency were assessed and recorded.

Result and discussion

Effect of spawn density on the sporophore yield of *Lentinus edodes*

Data pertaining to evaluation of optimum spawn density (Table 1) indicated that the spawn density of 20 per cent level completely colonized the beds within 31.3 days and the same treatment recorded comparatively less number of days for browning and pin head formation. However an increase in the spawn density beyond 20 per cent level resulted in reduced yield. The spawn density of ten and five per cent levels recorded significantly lesser yield. The fast mycelial growth observed in 20 per cent level of spawn density could be due to rapid colonization of the substrate which minimized the chances of contamination by the competitors and thus attributed to higher yields. Many workers have reported beneficial effect of increased rate of spawning (Pandey *et al.*, 2000; Nwanze *et al.*, 2005). However in the present study, it was also observed that the increase in the spawn density beyond 20 per cent resulted in decreased yield. This might be due to the reason that the higher spawn density may have resulted in increase in temp., CO₂ conc. and decrease in the oxygen conc. of the substrate (Chitra arya and Arun arya , 2003).

Effect of bed substrate on the growth and yield of *Lentinus edodes*

The present study was aimed to find out the suitability of various commonly available agro and agro-industrial wastes as substrates for shiitake cultivation. The results on the influence of various substrates are presented in table 2. The substrates, silver oak saw dust recorded shortest spawn run (32.5 days), browning period (57.5 days), sporophore yield (340.10g) with biological efficiency of 68 per cent followed by eucalyptus saw dust. The wood (natural) log cultivation cycle usually takes six years with low biological efficiency (less than 33 per cent). In the last decades, indoor shiitake production has been introduced substituting wood logs by so called synthetic media, sawdust being the basic ingredient for substrate formula (Pryzbylowicz and Donoghue, 1998).The productivity of this system depends on the quality of the sawdust used as substrate (Song *et al.*, 1989). Sawdust is difficult to standardize because it is generally obtained from residues of different types of



wood. Thus, the availability of low-cost and homogeneous substrate for production is an important factor for successful cultivation. However, a widely used standard substrate formula is 80 per cent hardwood sawdust and 20 per cent supplements on a dry weight basis (Royse 1985; Miller and Jong, 1987). Some formulation, with all ingredients based on oven dry substrate weight, consisting of 80 per cent sawdust and 20 per cent bran in Asia (Lizuka and Takeuchi, 1978); 80 per cent sawdust , 10 per cent bran and 10 per cent wheat or millet in USA (Miller and Jong, 1987) and 84 per cent sawdust, 5 per cent rice bran, 5 per cent wheat bran, 3 per cent soybean and 3 per cent lime in Taiwan (Kalberer, 1987), are commonly used for *L. edodes* cultivation as standard substrates. The present findings are also in line with the abovefindings.



Table 1: Effect of spawn density on the sporophore yield of *L.edodes*

Tr.no	Spawn density (g/bed)	Spawn run (days)	Browning (days)	Pin head formation (days)	Soprophore (wt)	Sporophore number/bed	Yield (g/bed)	Biological efficiency (%)
1	5	36.8 e	74.8f	96.50e	15.3f	14.1e	216f	43.2e
2	10	35.4 d	72.3e	92.00d	16.4e	17.5a	287e	57.4d
3	15	32.9b	71.0d	88.0c	18.3d	16.9b	309d	61.8c
4	20	31.3a	57.8a	83.1a	21.1a	15.9d	335a	67.0a
5	25	32.5 b	60.7b	85.3b	20.2b	16.0c	323b	64.6b
6	30	33.1c	64.3c	87.2c	19.1c	16.7b	318c	63.6b



Sl.No.	Substrates	Spawn run (days)	Browning (days)	Days for first harvest	Total yield (g/kg)	B.E %
1	Teak saw dust	35.2 _c	65.2 _c	104.3 _c	296.25 _c	59.2 _c
2	Silver oak saw dust	32.5 _a	57.5 _a	95.2 _a	340.10 _a	68.0 _a
3	Groundnut shell	37.9 _d	71.2 _d	112.0 _d	279.80 _d	55.7 _e
4	Coconut saw dust	37.0 _d	70.9 _d	110.5 _d	287.75 _d	57.5 _d
5	Coir pith	41.5 _g	80.3 _h	134.8 _h	260.0 _f	52.0 _g
6	Eucalyptus saw dust	33.4 _b	61.2 _b	98.5 _b	310.20 _b	62.0 _b
7	Sugarcane trash	39.7 _e	76.3 _f	126.8 _f	269.50 _e	53.9 _f
8	Sugarcane baggase	38.3 _d	74.4 _e	122.7 _e	273.40 _e	54.6 _f
9	Paddy straw	40.2 _f	78.1 _g	130.3 _g	250.70 _g	50.1 _h

Table 2: Effect of bed substrate on the growth and yield of *L.edodes*



Selected References

1. Sharma, S.R., Kumar, S. and Sharma, V.P. 2006. Physiological Requirement for Cultivation of Malaysian Strain of Shiitake, *Lentinula edodes*. *J. Mycol. PL. Pathol.*, 36(2):149-152
2. Sivaprakasam, K. 1980. Studies on oyster mushroom *Pleurotus sajor-caju* (Fr.) Singer. Ph.D. Thesis, Tamil Nadu Agricultural University, India.
3. Thiribhuvanamala, G., Krishnamoorthy, A.S., Shanthi, K. and Marimuthu, T. 2005a. Development of suitable substrates for spawn and mushroom production of *Lentinula edodes* and *Auricularia polytricha*. *Madras Agric. J.*, 92(4-6):344-348.
4. Royse, D.J., 2005. Cultivation of shiitake on natural and synthetic logs. Penn State's College of Agricultural Science. <http://www.cas.psu.edu/FreePubs/pdfs/ul203>.
5. Pandey, A., Soccol, C.R., Nigam, P., Soccol, V.T., Vandenerghe, L.P.S. and Monhan, R. 2000. Biotechnological potential of agroindustrial residues II: cassava bagasse. *Bioresource Technology*, 74:81-87. Pandey, M. 2005. Mushroom Research and Development of India. <http://www.mushworld.com/oversea/view.asp?id=5140&vid=7165>
6. Nwanze, P.I., Khan, A.U., Ameh, J.B. and Umoh, V.J. 2005. The effect of spawn grains, culture media, oil types and rates on carpophore production of *Lentinus squarrosulus* (Mont.) Singer. *Afr. J. Biotechnol.*, 4(6): 472-477.
7. Chitra Arya and Arun Arya. 2003. Effect of acid hydrolysis of substrate on the yield of oyster mushroom (*Pleurotus sajor-caju* (Fr.) Singer). *Mushroom Res.*, 12(1): 35-38.
8. Przybylowicz, P. and Donoghue, J. 1988. Shiitake growers' handbook. Kendall/Hunt Publishing Co., Dubuque, Iowa, pp.457.
9. Miller, M.W. and Jong, S.C. 1987. Commercial cultivation of shiitake in sawdust filled plastic bags. In: Wuest, P.J., Royse, D.J., Beelman, R.B. (Eds.): *Development in Crop Science, Cultivating Edible Fungi*, Elsevier Publishers, The Netherlands, pp.421-426.
10. Kalberer, P.P. 1987. Experiments on the cultivation of shiitake (*Lentinus edodes*) on sawdust. *Mushroom Sci.*, 12:231-235.